

PRODUCT INFORMATION

| | |
|------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Clone ID | DMC686 |
| Target | ALPP |
| Synonyms | ALP; ALPI; IAP; PALP; PLAP; PLAP-1 |
| Host Species | Rabbit |
| Description | Anti-ALPP antibody(DMC686); IgG1 Chimeric mAb |
| Delivery | In Stock |
| Uniprot ID | P05187 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is an alkaline phosphatase; a metalloenzyme that catalyzes the hydrolysis of phosphoric acid monoesters. It belongs to a multigene family composed of four alkaline phosphatase isoenzymes. The enzyme functions as a homodimer and has a catalytic site containing one magnesium and two zinc ions; which are required for its enzymatic function. One of the main sources of this enzyme is the liver; and thus; it's one of several indicators of liver injury in different clinical conditions. In pregnant women; this protein is primarily expressed in placental and endometrial tissue; however; strong ectopic expression has been detected in ovarian adenocarcinoma; serous cystadenocarcinoma; and other ovarian cancer cells. [provided by RefSeq; Aug 2020] |
| Usage | Research use only |
| Conjugate | Unconjugated |
| DIMA Disclaimer | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. |






Figure 1. Flow cytometry analysis with Anti-ALPP(DMC686) on HEK293 cells transfected with human ALPP(Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

