

PRODUCT INFORMATION

Clone ID	DM53
Target	B7-H3
Synonyms	B7-H3; CD276; B7 homolog 3; B7H3
Host Species	Rabbit
Description	Anti-B7-H3 antibody(DM53); Rabbit mAb
Delivery	In Stock
Uniprot ID	Q5ZPR3
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene belongs to the immunoglobulin superfamily; and thought to participate in the regulation of T-cell-mediated immune response. Studies show that while the transcript of this gene is ubiquitously expressed in normal tissues and solid tumors; the protein is preferentially expressed only in tumor tissues. Additionally; it was observed that the 3' UTR of this transcript contains a target site for miR29 microRNA; and there is an inverse correlation between the expression of this protein and miR29 levels; suggesting regulation of expression of this gene product by miR29. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr







Figure 1. Flow cytometry data of serially titrated Rabbit anti-B7H3 monoclonal antibody (**clone: DM53**) on HEK293 cell line transfected with human B7-H3. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

A



B




Figure 2. HEK293 cell line transfected with irrelevant protein **(A)** and human B7-H3 **(B)** were surface stained with Rabbit anti-B7-H3 monoclonal antibody 1μg/ml (**clone: DM53**) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.

