

PRODUCT INFORMATION

Clone ID	DM172
Target	CD114
Synonyms	CSF3R;CD114;GCSFR
Host Species	Rabbit
Description	Anti-CD114 antibody(DM172); Rabbit mAb
Delivery	In Stock
Uniprot ID	Q99062
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is the receptor for colony stimulating factor 3; a cytokine that controls the production; differentiation; and function of granulocytes. The encoded protein; which is a member of the family of cytokine receptors; may also function in some cell surface adhesion or recognition processes. Alternatively spliced transcript variants have been described. Mutations in this gene are a cause of Kostmann syndrome; also known as severe congenital neutropenia.
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. Flow cytometry analysis with Anti-CD114 (DM172) on HEK293 cells transfected with human CD114 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of rabbit anti-human CD114 mAb(DME100172).
(A) DME100172 does not bind to CHO-S cells that do not express CD114.
(B) A clear peak shift of DME100172 was seen compared to the control when incubated with CD114-expressing THP-1 cells, indicating strong binding of DME100172 to CD114. Antibodies were incubated at 5 µg/mL.

