

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Clone ID | 1A3 |
| Target | CD14 |
| Synonyms | CD14 Molecule;Myeloid Cell-Specific Leucine-Rich Glycoprotein;CD14 Antigen;Monocyte Differentiation Antigen CD14 |
| Host Species | Rabbit |
| Description | Anti-CD14 antibody(1A3), IgG1 Chimeric mAb |
| Delivery | In Stock |
| Uniprot ID | P08571 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1/100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide, and to viruses. This gene has been identified as a target candidate in the treatment of SARS-CoV-2-infected patients to potentially lessen or inhibit a severe inflammatory response. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Aug 2020] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Flow cytometry analysis with 10 μ g/mL Anti-CD14 (1A3) mAb on HEK293 cells transfected with human CD14 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

