

## PRODUCT INFORMATION

|   |   |
|---|---|
| <b>Clone ID</b>                         | 2D2   |
| <b>Target</b>                           | CD147   |
| <b>Synonyms</b>                         | Basigin; BSG; 5F7; CD147; EMMPRIN; M6; OK; TCSF   |
| <b>Host Species</b>                     | Rabbit  |
| <b>Description</b>                      | Anti-CD147 antibody(2D2), IgG1 Chimeric mAb   |
| <b>Delivery</b>                         | In Stock  |
| <b>Uniprot ID</b>                       | P35613  |
| <b>IgG type</b>                         | Rabbit/Human Fc chimeric IgG1   |
| <b>Clonality</b>                        | Monoclonal  |
| <b>Reactivity</b>                       | Human   |
| <b>Applications</b>                     | Flow Cyt  |
| <b>Recommended Dilutions</b>            | Flow Cyt 1/100  |
| <b>Purification</b>                     | Purified from cell culture supernatant by affinity chromatography   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | The protein encoded by this gene, basigin, is a plasma membrane protein that is important in spermatogenesis, embryo implantation, neural network formation, and tumor progression. Basigin is also a member of the immunoglobulin superfamily, ubiquitously expressed in various tissues. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2020] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |
| <b>DIMA Disclaimer</b>                  | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr  |






Figure 1. Flow cytometry analysis with 1 $\mu$ g/mL Anti-CD147 (2D2) mAb on HEK293 cells transfected with human CD147 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human CD147 mAb(DMC101093).

(A) DMC101093 does not bind to CHO-S cells that do not express CD147.

(B) A clear peak shift of DMC101093 was seen compared to the control when incubated with CD147-expressing THP-1 cells, indicating strong binding of DMC101093 to CD147. Antibodies were incubated at 5  $\mu$ g/ml.

