

## PRODUCT INFORMATION

<b>Clone ID</b>	DM146
<b>Target</b>	CD155
<b>Synonyms</b>	PVR; FLJ25946; PVS; CD155; TAGE4; HVED; NECL5
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-CD155 antibody(DM146); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P15151
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a transmembrane glycoprotein belonging to the immunoglobulin superfamily. The external domain mediates cell attachment to the extracellular matrix molecule vitronectin; while its intracellular domain interacts with the dynein light chain Tctex-1:DYNLT1. The gene is specific to the primate lineage; and serves as a cellular receptor for poliovirus in the first step of poliovirus replication. Multiple transcript variants encoding different isoforms have been found for this gene.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr






Figure 1. ELISA plate pre-coated by 1  $\mu$ g/ml (100  $\mu$ l/well) Human CD155 protein, mFc tagged protein PME100027 can bind Rabbit anti-CD155 monoclonal antibody (clone: DM146) in a linear range of 5-100 ng/ml.




Figure 2. CD155 protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with Anti-CD155 (DM146) on HEK293 cells transfected with human CD155 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).






Figure 3. Flow cytometry analysis of antigen binding of rabbit anti-human CD155 mAb(DME100146).  
(A) DME100146 does not bind to CHO-S cells that do not express CD155.  
(B) A clear peak shift of DME100146 was seen compared to the control when incubated with CD155-expressing Daudi cells, indicating strong binding of DME100146 to CD155. Antibodies were incubated at 5  $\mu$ g/ml.

