

PRODUCT INFORMATION

Clone ID **DMC280** CD21 **Target**

C3DR; CD21; CR; CVID7; SLEB9 **Synonyms**

Host Species Rabbit

Description Anti-CD21 antibody(DMC280); IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** P20023

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human

Applications Flow Cyt; WB

Recommended **Dilutions**

Flow Cyt 1:100; WB 1:1000

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Background

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a membrane protein; which functions as a receptor for Epstein-Barr virus (EBV) binding on B and T lymphocytes. Genetic variations in this gene are associated with

susceptibility to systemic lupus erythematosus type 9 (SLEB9). Alternatively spliced transcript variants encoding different isoforms have been

found for this gene.


Usage Research use only

Conjugate Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or **DIMA Disclaimer**

reverse engineering attempt is prohibited. We are

actively scr



Email: info@dimabio.com Website: www.dimabio.com



Cat. No. DMC100280






Figure 1. Flow cytometry analysis with Anti-CD21 (DMC280) on HEK293 cells transfected with human CD21 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).



Figure 2.Anti-CD21 antibody (SKU# DMC100280) at 1/1000 dilution

Lane: RAJI, whole cell lysate

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 113 kDa AbdersecWaharsinetills@ kDa Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA)/+86-18062749453(China)/+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com









Figure 3. Flow cytometry analysis of antigen binding of anti-human CD21 mAb(DMC100280). (A) DMC100280 does not bind to Jurkat cells that do not express CD21. (B) A clear peak shift of DMC100280 was seen compared to the control when incubated with CD21-expressing Raji cells, indicating strong binding of DMC100280 to CD21. Antibodies were incubated at $10 \, \mu g/mL$.



/+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

