

PRODUCT INFORMATION

Clone ID	DM204
Target	CD73
Synonyms	CD73;NT5E;5'-Nucleotidase;5'-NT;NT5;NTE
Host Species	Rabbit
Description	Anti-CD73 antibody(DM204); Rabbit mAb
Delivery	In Stock
Uniprot ID	P21589
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a plasma membrane protein that catalyzes the conversion of extracellular nucleotides to membrane-permeable nucleosides. The encoded protein is used as a determinant of lymphocyte differentiation. Defects in this gene can lead to the calcification of joints and arteries. Two transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq; Mar 2011]
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. Flow cytometry analysis with Anti-CD73 (DM204) on HEK293 cells transfected with human CD73 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of Rabbit anti-human CD73 mAb(DME100204).
(A) DME100204 does not bind to Jurkat cells that do not express CD73.
(B) A clear peak shift of DME100204 was seen compared to the control when incubated with CD73-expressing HT55 cells, indicating strong binding of DME100204 to CD73. Antibodies were incubated at 10 μ g/mL.

