

## **PRODUCT INFORMATION**

Clone ID **DMC486 Target** CD23

**Synonyms** BLAST-2; CD23; CD23A; CLEC4J; FCE2; IGEBF

**Host Species** Rabbit

Anti-CDH23 antibody(DMC486); IgG1 Chimeric Description mAb

**Delivery** In Stock **Uniprot ID** P06734

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Storage & Shipping

**Background** 

**DIMA Disclaimer** 

Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

The protein encoded by this gene is a B-cell specific antigen; and a low-affinity receptor for IgE. It has essential roles in B cell growth and differentiation; and the regulation of IgE production. This protein also exists as a soluble secreted form; then functioning as a potent

mitogenic growth factor. Alternatively spliced transcript variants encoding different isoforms have been described for this gene [provided by

RefSeq; Jul 2011]

Usage Research use only Conjugate Unconjugated

All DIMA recombinant antibodies are genuinely

generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

> Email: info@dimabio.com Website: www.dimabio.com

actively scrutinizing all patent application to ensure no IP infringement.



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Flow cytometry analysis with Anti-CD23 (DMC486) on HEK293 cells transfected with human CD23 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).





Figure 2. Flow cytometry analysis of antigen binding of anti-human CD23 mAb(DMC100486). (A) DMC100486 does not bind to jurkat cells that do not express CD23. (B) A clear peak shift of DMC100486 was seen compared to the control when incubated with CD23-expressing Raji cells, indicating strong binding of DMC100486 to CD23. Antibodies were incubated at 5  $\mu$ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

