

PRODUCT INFORMATION

Common Name	AMG-757 without CD3
Conjugate	Unconjugated
Synonyms	SCDO1
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	DLL3
Uniprot ID	Q9NYJ7
Description	Anti-DLL3(tarlatamab without CD3 biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only

CONFIDENTIAL






Figure 1. Flow cytometry analysis with 15µg/mL Anti-DLL3(tarlatamab without CD3 biosimilar) mAb(BME100257) on 293T-DLL3 cells (Blue histogram) or isotype control mAb on 293T-DLL3 cells (Red histogram).

DIMABIO CONFIDENTIAL

