

PRODUCT INFORMATION

Clone ID	DM96
Target	DNAM1
Synonyms	DNAM1; CD226; PTA1
Host Species	Rabbit
Description	Anti-DNAM-1 antibody(DM96); Rabbit mAb
Delivery	In Stock
Uniprot ID	Q15762
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a glycoprotein expressed on the surface of NK cells; platelets; monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. The protein mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Alternative splicing results in multiple transcript variants.
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr






Figure 1. ELISA plate pre-coated by 2 $\mu\text{g/ml}$ (100 $\mu\text{l/well}$) Human DNAM-1 protein, mFc-His tagged protein ([getskuurl sku="PME100050"]) can bind Rabbit anti-DNAM-1 monoclonal antibody (**clone: DM96**) in a linear range of 0.64-80 ng/ml.




Figure 2. HEK293 cell line transfected with irrelevant protein (**A**) and human DNAM-1 (**B**) were surface stained with Rabbit anti-DNAM-1 monoclonal antibody 1 $\mu\text{g/ml}$ (**clone: DM96**) followed by PE-conjugated anti-rabbit IgG secondary antibody.

