

PRODUCT INFORMATION

Clone ID **DMC424 Target EREG**

Ep; EPR; ER **Synonyms Host Species** Rabbit

Description Anti-EREG antibody(DMC424); IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** 014944

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended **Dilutions**

Flow Cyt 1:100

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Background

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a secreted peptide hormone and member of the epidermal growth factor (EGF) family of proteins. The encoded protein is a ligand of the epidermal growth factor receptor (EGFR) and the structurally related erb-b2 receptor tyrosine kinase 4 (ERBB4). The encoded protein may be involved in a wide range of biological

processes including inflammation; wound healing; oocyte maturation; and cell proliferation. Additionally; the encoded protein may promote the progression of cancers of various human

tissues.


Usage Research use only Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing apparent application to

> > Email: info@dimabio.com Website: www.dimabio.com

DIMA Disclaimer

ensure no IP infringement.



/+86-400-006-0995(China)









Figure 1. Flow cytometry analysis with Anti-EREG (DMC424) on HEK293 cells transfected with human EREG (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).



Figure~2.~Flow~cytometry~analysis~of~antigen~binding~of~anti-human~EREG~mAb (DMC100424).

(A) DMC100424 does not bind to Jurkat cells that do not express EREG. (B) A clear peak shift of DMC100424 was seen compared to the control when incubated with EREG-expressing SiHa cells, indicating strong binding of DMC100424 to EREG. Antibodies were incubated at 5 μ g/mL.

Email: info@dimabio.com Website: www.dimabio.com

