Cat. No. DME100175



## **PRODUCT INFORMATION**

Clone ID DM175 **Target** HFR3

**Synonyms** HER3; ERBB3

**Host Species** Rabbit

Description Anti-Her3 antibody(DM175); Rabbit mAb

**Delivery** In Stock **Uniprot ID** P21860 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human

**Applications** ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Formulation & Reconstitution

Background

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However; it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and:or

overexpression of its protein have been reported in numerous cancers; including prostate; bladder; and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been

characterized. One isoform lacks the

intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported; but they have not been

thoroughly characterized.

Usage Research use only Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

> > Email: info@dimabio.com


Website: www.dimabio.com

**DIMA Disclaimer** reverse engineering attempt is prohibited. We are

actively scr

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)





**Figure 1.** Flow cytometry analysis with Anti-Her3 (DM175) on HEK293 cells transfected with human Her3 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

