Cat. No. BME100244



## **PRODUCT INFORMATION**

**Uniprot ID** P21860

**Common Name** MM-121, MM121, SAR256212

Conjugate Unconjugated HER3, ERBB3 **Synonyms Applications** ELISA, Flow Cyt

Recommended

ELISA 1:5000-10000, Flow Cyt 1:100 **Dilutions** 

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

**Host Species** Homo sapiens

Human IgG2(S378A) - Kappa IgG type

Reactivity Human **Target** HER3

**Description** Anti-Her3(seribantumab biosimilar) mAb

**Delivery** In Stock

Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping

at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient

> Email: info@dimabio.com Website: www.dimabio.com


témperature.

Research grade biosimilar. Not for use in

**Background** therapeutic or diagnostic procedures for humans

or animals.

Usage Research use only



/+86-400-006-0995(China)





## Anti-Her3(seribantumab biosimilar) mAb ELISA

0.2 μg of Human Her3, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human Her3 Protein, His Tag (PME100088) can bind Anti-Her3(seribantumab biosimilar) mAb (BME100244) in a linear range of 0.13–16 ng/mL.




Figure 2. Flow cytometry analysis with  $1\mu g/mL$  Anti-Her3(seribantumab biosimilar) mAb (BME100244) on HEK293 cells transfected with Human Her3 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com



Cat. No. BME100244






Figure 3. Flow cytometry analysis of antigen binding of anti-human Her3 mAb(BME100244). (A) BME100244 does not bind to 293T cells that do not express Her3. (B) A clear peak shift of BME100244 was seen compared to the control when incubated with Her3-expressing HT55 cells, indicating strong binding of BME100244 to Her3. Antibodies were incubated at 5  $\mu$ g/mL.



