

## **PRODUCT INFORMATION**

Clone ID 1G1 IL18BP **Target** 

**Synonyms** FVH:IL18BPa

**Host Species** Rabbit

Description Anti-IL18BP antibody(1G1), IgG1 Chimeric mAb

**Delivery** In Stock **Uniprot ID** 095998

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications ELISA** 

Recommended

**Background** 

ELISA 1/5000-10000 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene functions as an inhibitor of the proinflammatory cytokine, IL18. It binds IL18, prevents the binding of IL18 to its receptor, and thus inhibits IL18-induced IFNgamma production, resulting in reduced T-helper type 1 immune responses. This protein is

constitutively expressed and secreted in mononuclear cells. Elevated level of this protein is detected in the intestinal tissues of patients with Crohn's disease. Alternatively spliced transcript

variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011]


Usage Research use only Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under

patent application. Any protein sequencing or **DIMA Disclaimer** 

reverse engineering attempt is prohibited. We are

actively scr



Email: info@dimabio.com Website: www.dimabio.com





## Anti-IL18BP (1G1) mAb ELISA 0.1µg of IL18BP, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1  $\mu$ g/ml (100  $\mu$ l/well) Human IL18BP, hFc tagged protein can bind Rabbit anti-IL18BP monoclonal antibody(clone: 1G1) in a linear range of 5-100 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

