

PRODUCT INFORMATION

Common Name ACZ885

Conjugate Unconjugated **Synonyms** IL-1 beta;IL1F2 **Applications** ELISA; Flow Cyt

Recommended

Dilutions

ELISA 1:5000-10000; Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Host Species Homo sapiens

IgG type Human IgG1 - kappa

Reactivity Human **Target** IL1B **Uniprot ID** P01584

Description Anti-IL1B(canakinumab biosimilar) mAb

Delivery In Stock

Storage & Shipping

Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Email: info@dimabio.com Website: www.dimabio.com

Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

Background therapeutic or diagnostic procedures for humans

or animals.

Usage Research use only





Anti-IL1B (canakinumab biosimilar) mAb ELISA

0.1 µg of Human IL1B, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Human IL1B Protein, hFc Tag (PME100838) can bind Anti-IL1B Neutralizing antibody (BME100116) in a linear range of 0.61–156.25 ng/mL. In order to specifically detect BME100116, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis under cell membrane permeable condition with 1 μ g/mL Anti-IL1B (canakinumab biosimilar) mAb (BME100116) on HEK293 cells transfected with Human IL1B protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com