

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Clone ID | DMC272 |
| Target | IL7RA |
| Synonyms | CD127; CDW127; IL-7R-alpha; IL7RA; ILRA |
| Host Species | Rabbit |
| Description | Anti-IL7RA antibody(DMC272); IgG1 Chimeric mAb |
| Delivery | In Stock |
| Uniprot ID | P16871 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is a receptor for interleukin 7 (IL7). The function of this receptor requires the interleukin 2 receptor; gamma chain (IL2RG); which is a common gamma chain shared by the receptors of various cytokines; including interleukins 2; 4; 7; 9; and 15. This protein has been shown to play a critical role in V(D)J recombination during lymphocyte development. Defects in this gene may be associated with severe combined immunodeficiency (SCID). Alternatively spliced transcript variants have been found. |
| Usage | Research use only |
| Conjugate | Unconjugated |
| DIMA Disclaimer | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. |






Figure 1. Flow cytometry analysis with Anti-IL7RA (DMC272) on HEK293 cells transfected with human IL7RA (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

