

PRODUCT INFORMATION

Common Name	7F3com-2H2, MEDI-528
Conjugate	Unconjugated
Synonyms	Interleukin-9;IL-9;Cytokine P40;T-cell growth factor P40
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	IL9
Uniprot ID	P15248
Description	Anti-IL9(enokizumab biosimilar) mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Anti-IL9 (enokizumab biosimilar) mAb ELISA
0.1 µg of Cynomolgus IL9, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 µg/mL (100 µL/well) Cynomolgus IL9 Protein, hFc Tag (PME-C100003) can bind Anti-IL9 Neutralizing antibody (BME100122) in a linear range of 2.44-156.25 ng/mL. In order to specifically detect BME100122, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis under cell membrane permeable condition with 1 µg/mL Anti-IL9 (enokizumab biosimilar) mAb (BME100122) on HEK293 cells transfected with Human IL9 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

