

## **PRODUCT INFORMATION**

**Clone ID** 2H10 MC4R **Target** BMIQ20 **Synonyms Host Species** Rabbit

Description Anti-MC4R antibody(2H10), Rabbit mAb

**Delivery** In Stock **Uniprot ID** P32245 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended Flow Cyt 1/100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Storage & Shipping intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). The protein encoded by this gene is a membranebound receptor and member of the melanocortin receptor family. The encoded protein interacts with adrenocorticotropic and MSH hormones and is mediated by G proteins. This is an intronless gene. Defects in this gene are a cause of

autosomal dominant obesity. [provided by

RefSeq, Jan 2010]

**Usage** Research use only

Conjugate Unconjugated

**Background** 



Email: info@dimabio.com Website: www.dimabio.com






Figure 1. Flow cytometry analysis with  $2\mu g/mL$  Anti-MC4R (2H10) mAb on HEK293 cells transfected with human MC4R (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

