

## PRODUCT INFORMATION

Clone ID 8C4 MMAE **Target** N.A. **Synonyms Host Species** Rabbit

Description Anti-MMAE antibody(8C4); Rabbit mAb

**Delivery** In Stock **Uniprot ID** N.A.

IgG type Rabbit IgG Clonality Monoclonal

Reactivity N.A. **Applications ELISA** 

Recommended **Dilutions** 

ELISA 1:5000-10000

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

**Background** 


**DIMA Disclaimer** 

**Usage** Research use only Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

reverse engineering attempt is prohibited. We are

actively scr



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com





## ELISA assay to evaluate Anti-MMAE antibody 0.2μg Human IgG-MMAE per well




Figure 1. Elisa plates were pre-coated with IgG-MMAE (0.2 $\mu$ g/per well). Serial diluted anti-MMAE monoclonal antibody (DME101004) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-MMAE monoclonal antibody binding with IgG-MMAE is 1.356ng/ml.



