

PRODUCT INFORMATION

Common Name IMC RON-8, IMC-RON8, RON8

CD136;CDw136;NPCA3;PTK8;RON **Synonyms**

Conjugate Unconjugated

Applications ELISA

Recommended

ELISA 1:5000-10000 **Dilutions**

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Host Species Homo sapiens

IgG type Human IgG1 - kappa

Reactivity Human **Target** MST1R **Uniprot ID** Q04912

Anti-MST1R(narnatumab biosimilar) mAb **Description**

Delivery In Stock

Storage & Shipping

Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Email: info@dimabio.com Website: www.dimabio.com

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

Background therapeutic or diagnostic procedures for humans

or animals.

Usage Research use only





Anti-MST1R (narnatumab biosimilar) mAb ELISA

0.2 μg of Human MST1R, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human MST1R Protein, hFc Tag (PME100821) can bind Anti-MST1R Neutralizing antibody (BME100118) in a linear range of 0.73–46.88 ng/mL. In order to specifically detect BME100118, mouse anti-human Fab-specific antibody was used as detection antibody.

Email: info@dimabio.com Website: www.dimabio.com

