

PRODUCT INFORMATION

| | |
|------------------------------|---|
| Common Name | NP-137 |
| Conjugate | Unconjugated |
| Synonyms | NTN1; NTN1L |
| Applications | ELISA |
| Recommended Dilutions | ELISA 1:5000-10000 |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Host Species | Humanized |
| IgG type | Human IgG1 - kappa |
| Reactivity | Human |
| Target | Netrin-1 |
| Description | Anti-Netrin-1(NP-137 biosimilar) mAb |
| Delivery | In Stock |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature. |
| Background | Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. |
| Usage | Research use only |



Anti-Netrin-1(NP-137 biosimilar) mAb ELISA

0.2 μ g of Human Netrin-1, Flag tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human Netrin-1 Protein, Flag Tag (PME101766) can bind Anti-Netrin-1(NP-137 biosimilar) mAb (BME100199) in a linear range of 0.64–16 ng/mL.

