

PRODUCT INFORMATION

Uniprot ID	P26718
Common Name	5C5
Conjugate	Unconjugated
Synonyms	NKG2D,CD314,KLRK1,NK cell receptor D
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	NKG2D
Description	Anti-NKG2D(5C5 biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only



Anti-NKG2D(5C5 biosimilar) mAb ELISA

0.2 µg of Human NKG2D, mFc tagged protein per well




Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human NKG2D Protein, mFc Tag (PME100079) can Anti-NKG2D(5C5 biosimilar) mAb (BME100207) in a linear range of 0.13-16 ng/mL.




Figure 2. Flow cytometry analysis with 1µg/mL Anti-NKG2D(5C5 biosimilar) mAb (BME100207) on HEK293 cells transfected with Human NKG2D and DAP10 protein (Blue histogram) or HEK293 transfected with DAP10 protein(Red histogram).



Anti-NKG2D(5C5 biosimilar) mAb ELISA

0.2 µg of Anti-NKG2D(5C5 biosimilar) mAb per well




Figure 3. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Anti-NKG2D(5C5 biosimilar) mAb (BME100207) can bind Biotinylated Human NKG2D Protein, hFc Tag (PME101956B) in a linear range of 200-1000 ng/mL.

