



## PRODUCT INFORMATION

<b>Clone ID</b>	DM112
<b>Target</b>	OX40 Ligand
<b>Synonyms</b>	OX40L; TNFSF4; CD252; Glycoprotein Gp34; TXGP1; CD134 ligand; CD134L
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-OX40 Ligand antibody(DM112); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P23510
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a cytokine of the tumor necrosis factor (TNF) ligand family. The encoded protein functions in T cell antigen-presenting cell (APC) interactions and mediates adhesion of activated T cells to endothelial cells. Polymorphisms in this gene have been associated with Sjogren's syndrome and systemic lupus erythematosus. Alternative splicing results in multiple transcript variants
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr





**Figure 1.** ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human OX40L protein, mFc-His tagged protein ([getskuurl sku="PME100021"]]) can bind Rabbit anti-OX40L monoclonal antibody (**clone: DM112**) in a linear range of 0.2-90 ng/ml.



**Figure 2.** Flow cytometry analysis with Anti-OX40L (**DM112**) on HEK293 cells transfected with human OX40L (Blue histogram) or HEK293 transfected with irrelevant protein(Red histogram).

