

PRODUCT INFORMATION

Common Name UC-961, cirmtuzumab

Conjugate Unconjugated ROR1;NTRKR1 **Synonyms Applications** ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions**

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution for specific instructions of reconstitution.

Host Species Humanized

IgG type Human IgG1 - kappa

Reactivity Human **Target** ROR1 **Uniprot ID** Q01973

Description Anti-ROR1 (zilovertamab biosimilar) mAb

Delivery In Stock

DIMA Disclaimer

Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in

Background therapeutic or diagnostic procedures for humans


or animals.

Usage Research use only

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

reverse engineering attempt is prohibited. We are

actively scr



Website: www.dimabio.com

Email: info@dimabio.com





Anti-ROR1 (zilovertamab biosimilar) mAb ELISA

0.1 μg of Human ROR1, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human ROR1, His tagged protein PME100399 can bind Anti-ROR1 Neutralizing antibody (BME100073) in a linear range of 0.64-16 μ g/ml.




Figure 2. Flow cytometry analysis with Anti-ROR1 (zilovertamab biosimilar) mAb 15 μ g/mL on HEK293 cells transfected with Human ROR1 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).



Email: info@dimabio.com Website: www.dimabio.com








Figure 3. Flow cytometry analysis of antigen binding of anti-human ROR1 mAb(BME100073). (A) A clear peak shift of BME100073 was seen compared to the control when incubated with ROR1-expressing 8226 cells, indicating strong binding of BME100073 to ROR1. Antibodies were incubated at 2 μ g/mL.



