

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | DM26 |
| Target | S protein RBD |
| Synonyms | SARS-CoV-2 RBD |
| Host Species | Rabbit |
| Description | Anti-SARS-CoV-2 RBD antibody(DM26); Rabbit mAb |
| Delivery | In Stock |
| Uniprot ID | P0DTC2 |
| IgG type | Rabbit IgG |
| Clonality | Monoclonal |
| Reactivity | SARS-CoV-2 |
| Applications | ELISA; Flow Cyt |
| Recommended Dilutions | ELISA 1:5000-10000; Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Endotoxin | Less than 1.0 EU/μg by the LAL method. For <1 EU/mg requirements, please contact us for customization. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits; S1 and S2. S1 mainly contains a receptor binding domain (RBD); which accounts for recognizing the cell surface receptor; ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response. |
| Usage | Research use only |
| Conjugate | Unconjugated |
| DIMA Disclaimer | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr |






Figure 1. Competition flow cytometry assay demonstrating Rabbit anti-RBD monoclonal antibody (**clone: DM26**) blockade of SARS-CoV-2 (COVID-19) S protein RBD (1 μ g/ml, [getskuurl sku="PME100497"]) binding to HEK293 cell line transfected with human ACE2. IC₅₀=1717ng/ml. The Y-axis represents the geometric mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

