

## **PRODUCT INFORMATION**

Clone ID **DM26** 

**Target** S protein RBD SARS-CoV-2 RBD **Synonyms** 

**Host Species** Rabbit

Anti-SARS-CoV-2 RBD antibody(DM26); Rabbit Description mAb

**Delivery** In Stock **Uniprot ID** P0DTC2 Rabbit IgG IgG type Clonality Monoclonal Reactivity SARS-CoV-2

**Applications** ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses


**Background** 

ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits; S1 and S2. S1 mainly contains a receptor binding domain (RBD); which accounts for recognizing the cell surface receptor; ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell

> Email: info@dimabio.com Website: www.dimabio.com


response.

Research use only **Usage** 









**Figure 1.** Competition flow cytometry assay demonstrating Rabbit anti-RBD monoclonal antibody (**clone: DM26**) blockade of SARS-CoV-2 (COVID-19) S protein RBD ( $1\mu$ g/ml, [getskuurl sku="PME100497"]) binding to Expi 293 cell line transfected with human ACE2. IC50=1717ng/ml. The Y-axis represents the geometric mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

Email: info@dimabio.com Website: www.dimabio.com

