

PRODUCT INFORMATION

Clone ID DMC392 B4GALT1 **Target**

Synonyms GGTB2; Beta4Gal-T1; b4Gal-T1; Nal synthase

Host Species

Biotinylated Anti-B4GALT1 antibody(DMC392); **Description**

IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** P15291

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **ELISA Applications**

Recommended

ELISA 1:5000-10000 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

> Email: info@dimabio.com Website: www.dimabio.com

témperature.



Cat. No. DMC100392B



This gene is one of seven beta-1,4galactosyltransferase (beta4GalT) genes. They encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose; all transfer galactose in a beta1,4 linkage to similar acceptor sugars: GlcNAc; Glc; and Xyl. Each beta4GalT has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins; they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a transmembrane anchor. By sequence similarity; the beta4GalTs form four groups: beta4GalT1 and beta4GalT2; beta4GalT3 and beta4GalT4; beta4GalT5 and beta4GalT6; and beta4GalT7. This gene is unique among the beta4GalT genes because it encodes an enzyme that participates both in glycoconjugate and lactose biosynthesis. For the first activity; the enzyme adds galactose to N-acetylglucosamine residues that are either monosaccharides or the nonreducing ends of glycoprotein carbohydrate chains. The second activity is restricted to lactating mammary tissues where the enzyme forms a heterodimer with alpha-lactalbumin to catalyze UDP-galactose D-glucose UDP lactose. The two enzymatic forms result from alternate transcription initiation sites and post-translational processing. Two transcripts; which differ only at the 5' end; with approximate lengths of 4.1 kb and 3.9 kb encode

cleaved to form the soluble lactose synthase. **Usage** Research use only

Conjugate Biotinylated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to

Email: info@dimabio.com Website: www.dimabio.com

the same protein. The longer transcript encodes the type II membrane-bound; trans-Golgi resident protein involved in glycoconjugate biosynthesis. The shorter transcript encodes a protein which is

ensure no IP infringement.

Background

DIMA Disclaimer




Cat. No. DMC100392B



Biotinylated Anti-B4GALT1 antibody(DMC392); IgG1 Chimeric mAb ELISA

 $0.2~\mu g$ of Human B4GALT1, hFc tagged protein per well



Biotinylated Anti-B4GALT1 antibody(DMC392); IgG1 Chimeric mAb (ng/mL)

Email: info@dimabio.com Website: www.dimabio.com

Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human B4GALT1 Protein, hFc Tag (PME100703) can bind Biotinylated Anti-B4GALT1 antibody(DMC392); lgG1 Chimeric mAb (DMC100392B) in a linear range of 16–80 ng/mL.

