

PRODUCT INFORMATION

Clone ID	DMC371
Target	GPC3
Synonyms	DGSX; GTR2-2; MXR7; OCI-5; SDYS; SGB; SGBS; SGBS1
Host Species	Rabbit
Description	Biotinylated Anti-GPC3 antibody(DMC371); IgG1 Chimeric mAb
Delivery	2-3 weeks
Uniprot ID	P51654
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPs) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation. The protein encoded by this gene can bind to and inhibit the dipeptidyl peptidase activity of CD26; and it can induce apoptosis in certain cell types. Deletion mutations in this gene are associated with Simpson-Golabi-Behmel syndrome; also known as Simpson dysmorphia syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq; Sep 2009] References [1] Fu Ying,Urbaniak Daniel J,Nani Roger R et al. Glypican-3-Specific Antibody Drug Conjugates Targeting Hepatocellular Carcinoma.[J].Hepatology; 2019; 70: 563-576. Zhang Yi-Fan,Horvath Mitchell,Humanization of high-affinity antibodies targeting glypican-3 in hepatocellular carcinoma.
Usage	Research use only
Conjugate	Biotinylated

DIMA Disclaimer

Address: Wuhan institute of Biotechnology B7, Biolake No.666
Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com



DIMABIO CONFIDENTIAL

