Cat. No. BME100212B



PRODUCT INFORMATION

Uniprot ID N/A

Common Name Hen egg Lysozyme

Conjugate Biotinylated

Synonyms HEL

Applications ELISA, Flow Cyt

Recommended

ELISA 1:5000-10000, Flow Cyt 1:100 **Dilutions**

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Host Species Chimeric IgG type IgG4 Reactivity N/A **Target** HFI

Biotinylated Anti-HEL Human IgG4-Kappa Isotype Description

control mAb

Delivery

Storage & Shipping

Background

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized antibodies are shipped at ambient

témperature.

Anti Chicken Hen Egg Lysozyme, specifically recognises Hen Egg Lysozyme (HEL), known also as muramidaes or N-acetylmuramidae

glycanhydrolase, a 14kDa enzymic protein involved in the destruction of bacteria. Lysozyme damages bacterial cell walls by catalyzing hydrolysis of 1,4-beta-linkages between Nacetylmuramic acid and N-acetyl-D-glucosamine residues in a peptidoglycan and between N-

acetyl-D-glucosamine residues in chitodextrins. Lysozyme is abundant in a number of secretions, such as tears, saliva, human milk and mucus. It is also present in cytoplasmic granules of PMN's and high concentrations of lysozyme are present in egg white. C-type lysozymes are closely related

to alpha-lactalbumin in sequence and structure making them part of the same family.

Usage Research use only

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

DIMA Disclaimer reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.





