

## **PRODUCT INFORMATION**

Clone ID DMC271 **Target** II<sub>1B</sub>

**Synonyms** IL-1; IL1-BETA; IL1F2

**Host Species** 

Biotinylated Anti-IL1B antibody(DMC271); IgG1 Description

Chimeric mAb

**Delivery** 2-3 weeks **Uniprot ID** P01584

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human **Applications** Flow Cyt

Recommended

Storage & Shipping

Background

Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a

proprotein; which is proteolytically processed to its active form by caspase 1 (CASP1:ICE). This cytokine is an important mediator of the inflammatory response; and is involved in a variety of cellular activities; including cell proliferation; differentiation; and apoptosis. The induction of cyclooxygenase-2 (PTGS2:COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly; IL-1B has been implicated in human estecarthritis nathogenesis

implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019 (COVID-19) present elevated levels of proinflammatory cytokines such as IL-1B in bronchial

alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent; a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family

> Email: info@dimabio.com Website: www.dimabio.com

genes form a cytokine gene cluster on chromosome 2

Usage Research use only

Conjugate Biotinylated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)



Biotinylated Anti-IL1B antibody(DMC271); IgG1 Chimeric mAb

Cat. No. DMC100271B



**DIMA Disclaimer** 

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to JIMABIO COLIFINITATION CONTRACTOR ensure no IP infringement.

> Email: info@dimabio.com Website: www.dimabio.com



