

PRODUCT INFORMATION

Target	BTN3A1
Synonyms	Butyrophilin protein
Description	Recombinant Cynomolgus BTN3A1 protein with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	A0A330KVC6
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	BTN3A1(Gln1-Ser218) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 24.3 kDa after removal of the signal peptide. The apparent molecular mass of cBTN3A1-His is approximately 25-35 kDa due to glycosylation.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Butyrophilin subfamily 3 member A1 (BTN3A1) is also known as CD277 and BTF5, which belongs to the immunoglobulin superfamily and contains one B30.2/SPRY domain and two Ig-like V-type (immunoglobulin-like) domains. BTN3A1 plays a role in T-cell activation and in the adaptive immune response. Also, BTN3A1 regulates the proliferation of activated T-cells and the release of cytokines and IFNG by activated T-cells. Furthermore, BTN3A1 mediates the response of T-cells toward infected and transformed cells that are characterized by high levels of phosphorylated metabolites, such as isopentenyl pyrophosphate.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Cynomolgus BTN3A1 Protein, His Tag on SDS-PAGE under reducing condition.

