Background



PRODUCT INFORMATION

BTLA Target

Monoclonal Cell Line Derived from K562 Cells Description Engineered for Stable Expression of Human BTLA

Using Lentiviral Technology

Host Cells K562 **Uniprot ID** Q7Z6A9 **Applications FACS Data**

RPMI-1640+10% FBS+1% P.S+1% Gln+2 ug/mL **Growth media**

Puromycin

Package 5E6 Cells/mL

Suggested Control SKU: BME100054

1. Please inspect cells upon receipt and report any issues promptly. 2. We offer one-time

replacements for issues reported within a week of

Warranty and receipt. 3. User-induced issues are not eligible for Disclaimer free replacements. 4. We do not accept liability for damages resulting from cell use, storage, or loss. 5. Feedback received more than one month

after receipt will not be processed.

Cells are shipped using dry ice and require liquid Storage & Shipping nitrogen storage for long term preservation.

Synonyms BTLA; CD272

> This gene encodes a member of the immunoglobulin superfamily. The encoded protein contains a single immunoglobulin (Ig) domain and is a receptor that relays inhibitory signals to suppress the immune response.

> Alternative splicing results in multiple transcript variants. Polymorphisms in this gene have been associated with an increased risk of rheumatoid

> > Email: info@dimabio.com Website: www.dimabio.com

arthritis.

Usage For research use only.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)





Hu_BTLA K562 Cell Line




Figure 1. Flow cytometry analysis of human BTLA overexpression using Hu_BTLA K562 Cell Line (Cat. No. CEL100091) and Anti-BTLA (icatolimab biosimilar)mAb (Cat. No. BME100054)

Email: info@dimabio.com Website: www.dimabio.com

