Warranty and Disclaimer

Background



PRODUCT INFORMATION

CCR5 **Target**

Monoclonal Cell Line Derived from K562 Cells, Engineered for Stable Expression of Human CCR5 Using Lentiviral Technology **Description**

Host Cells **Uniprot ID** P51681 **Applications FACS Data**

Growth media RPMI-1640+10% FBS+1% P.S+1% Gln+2 ug/mL Puromycin

5F6 Cells/ml Package **Suggested Control** SKU: BME100204

1. Please inspect cells upon receipt and report any issues promptly. 2. We offer one-time replacements for issues reported within a week of receipt. 3. User-induced issues are not eligible for free replacements. 4. We do not accept liability for damages resulting from cell use, storage, or loss. 5.

Feedback received more than one month after receipt will not

be processed.

Cells are shipped using dry ice and require liquid nitrogen Storage & Shipping

storage for long term preservation.


CC-CKR-5;CCCKR5;CCR-5;CD195;CKR-5;CKR5;CMKBR5;IDDM22 Synonyms

> This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. This protein is expressed by T cells and macrophages, and is known to be an important co-receptor for macrophage-tropic virus, including HIV, to enter host cells. Defective alleles of this gene have been associated with the HIV infection resistance. The ligands of this recenter include managery to champattractant protein 2 of this receptor include monocyte chemoattractant protein 2

> of this receptor include monocyte chemoattractant protein 2 (MCP-2), macrophage inflammatory protein 1 alpha (MIP-1 alpha), macrophage inflammatory protein 1 beta (MIP-1 beta) and regulated on activation normal T expressed and secreted protein (RANTES). Expression of this gene was also detected in a promyeloblastic cell line, suggesting that this protein may play a role in granulocyte lineage proliferation and differentiation. This gene is located at the chemokine receptor gene cluster region. An allelic polymorphism in this gene results in both functional and non-functional alleles; the reference genome represents the functional allele. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2015]

Email: info@dimabio.com Website: www.dimabio.com

Usage For research use only.










Figure 1. Flow cytometry analysis of human CCR5 overexpression using Hu_CCR5 K562 Cell Line (Cat. No. CEL100090) and Anti-CCR5(leronlimab biosimilar) mAb (Cat. No. BME100204)

Email: info@dimabio.com Website: www.dimabio.com

