

PRODUCT INFORMATION

Target	TACI
Description	Monoclonal Cell Line Derived from 293T Cells, Engineered for Stable Expression of Human TACI Using Lentiviral Technology
Host Cells	293T
Uniprot ID	O14836
Applications	FACS Data
Growth media	DMEM+10% FBS+1% P.S+Gln+2 ug/mL Puromycin
Package	5E6 Cells/mL
Suggested Control	SKU: IgG ch7B10
Warranty and Disclaimer	1. Please inspect cells upon receipt and report any issues promptly. 2. We offer one-time replacements for issues reported within a week of receipt. 3. User-induced issues are not eligible for free replacements. 4. We do not accept liability for damages resulting from cell use, storage, or loss. 5. Feedback received more than one month after receipt will not be processed.
Storage&Shipping	Cells are shipped using dry ice and require liquid nitrogen storage for long term preservation.
Synonyms	CD267;TACI;TNFRSF13B
Background	The protein encoded by this gene is a lymphocyte-specific member of the tumor necrosis factor (TNF) receptor superfamily. It interacts with calcium-modulator and cyclophilin ligand (CAML). The protein induces activation of the transcription factors NFAT, AP1, and NF-kappa-B and plays a crucial role in humoral immunity by interacting with a TNF ligand. This gene is located within the Smith-Magenis syndrome region on chromosome 17.
Usage	For research use only.



Hu_TACI 293T Cell Line




Figure 1. Flow cytometry analysis of human TACI overexpression using Hu_TACI 293T Cell Line (Cat. No. CEL100113) and Anti-TACI antibody(7B10); IgG1 Chimeric mAb (Cat. No. IgG ch7B10)

