

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	5HT1D
Synonyms	5-HT1D, HT1DA, HTR1DA, HTRL, RDC4
Description	Human 5HT1D full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P28221
Expression Host	HEK293
Protein Families	GPCR, Transmembrane, Druggable Genome, GPCRDB Class A Rhodopsin-like, Monoamine GPCRs,
Protein Pathways	
Molecular Weight	The human full length 5HT1D protein has a MW of 41.9kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for ergot alkaloid derivatives, various anxiolytic and antidepressant drugs and other psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylyl cyclase. Signaling inhibits adenylyl cyclase activity. Regulates the release of 5-hydroxytryptamine in the brain, and thereby affects neural activity. May also play a role in regulating the release of other neurotransmitters. May play a role in vasoconstriction.[UniProtKB/Swiss-Prot Function]
Usage	Research use only
Conjugate	Unconjugated

