

PRODUCT INFORMATION

|                              |  |
|------------------------------|--|
| Tag                          | C-Flag Tag   |
| Target                       | 5HT1E  |
| Synonyms                     | 5-HT1E   |
| Description                  | Human 5HT1E full length protein-synthetic nanodisc   |
| Delivery                     | 6~8weeks   |
| Uniprot ID                   | P28566   |
| Expression Host              | HEK293   |
| Protein Families             | GPCR,Transmembrane,Druggable Genome,   |
| Protein Pathways             | GPCRDB Class A Rhodopsin-like,Monoamine GPCRs,   |
| Molecular Weight             | The human full length 5HT1E protein has a MW of 41.7kDa  |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for  |
| Storage & Shipping           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| Background                   | G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various alkaloids and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling inhibits adenylate cyclase activity.[UniProtKB/Swiss-Prot Function] |
| Usage                        | Research use only  |
| Conjugate                    | Unconjugated   |

