

## **PRODUCT INFORMATION**

C-Flag&Strep Tag Tag

**Target** 5HT6R

**Synonyms** 5-HT6; 5-HT6R

Human 5HT6R-Strep full length protein-synthetic **Description** 

nanodisc In Stock

**Delivery Uniprot ID** P50406 **Expression Host HEK293** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

Calcium signaling pathway, Neuroactive ligand-**Protein Pathways** 

receptor interaction

The human full length 5HT6R-Strep protein has a **Molecular Weight** 

MW of 47.0 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

Formulation & Reconstitution for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater

than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

A protein that belongs to the seventransmembrane G protein-coupled receptor family of proteins. The encoded protein couples with the Gs alpha subunit and stimulates adenylate cyclase to activate the cyclic AMP-

dependent signaling pathway. This receptor is thought to regulate cholinergic neuronal

transmission in the brain. Several antidepressants and antipsychotic drugs have a high affinity for


this receptor.

Usage Research use only

Conjugate Unconjugated

**Background** 









## ELISA assay to evaluate 5HT6R-Strep-Nanodisc 0.2µg Human 5HT6R-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag 5HT6R-Strep-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with 5HT6R-Strep-nanodisc is 2.122ng/ml.



Figure 2. Human 5HT6R-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE



Email: info@dimabio.com Website: www.dimabio.com

