

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	ACKR1
Synonyms	CCBP1; CD234; DARC; DARC/ACKR1; Dfy; FY; GPD; GpFy; WBCQ1
Description	Human ACKR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	Q16570
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length ACKR1 protein has a MW of 35.6 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a glycosylated membrane protein and a non-specific receptor for several chemokines. The encoded protein is the receptor for the human malarial parasites Plasmodium vivax and Plasmodium knowlesi. Polymorphisms in this gene are the basis of the Duffy blood group system.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate ACKR1-Nanodisc 0.2 μ g Human ACKR1-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag ACKR1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ACKR1-Nanodisc is 37.84ng/ml.




Figure 2. Human ACKR1-Nanodisc, Flag Tag on SDS-PAGE

