

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CXCR7
Synonyms	ACKR3; CMKOR1; CXC-R7; CXCR-7; GPR159; RDC-1; RDC1
Description	Human CXCR7 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P25106
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length CXCR7 protein has a MW of 41.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A member of the G-protein coupled receptor family. Although this protein was earlier thought to be a receptor for vasoactive intestinal peptide (VIP), it is now considered to be an orphan receptor, in that its endogenous ligand has not been identified. The protein is also a coreceptor for human immunodeficiency viruses (HIV). Translocations involving this gene and HMGA2 on chromosome 12 have been observed in lipomas.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate CXCR7-Nanodisc
0.2 μ g Human CXCR7-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CXCR7-Nanodisc (0.2 μ g/per well). Serial diluted anti-CXCR7 monoclonal antibody (DMC100443) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CXCR7 monoclonal antibody binding with CXCR7-Nanodisc is 14.28ng/ml.




Figure2. Human CXCR7-Nanodisc, Flag Tag on SDS-PAGE

