

PRODUCT INFORMATION

ACVR2A **Target**

Synonyms ACVR2; ACTRII

Recombinant human ACVR2A Protein with C-Description

terminal human Fc tag

Delivery In Stock **Uniprot ID** P27037 **Expression Host HEK293**

Tag C-Human Fc tag

Molecular

Reconstitution

Background

Storage & Shipping

ACVR2A(Ala20-Pro135) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight**

39.5 kDa after removal of the signal peptide. The apparent molecular mass of ACVR2A-hFc is approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation &

 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a receptor that mediates the functions of activins, which are members of the transforming growth factor-beta (TGF-beta) superfamily involved in diverse biological processes. The encoded protein is a transmembrane serine-threonine kinase receptor

which mediates signaling by forming

heterodimeric complexes with various

combinations of type I and type II receptors and ligands in a cell-specific manner. The encoded type II receptor is primarily involved in ligand-binding and includes an extracellular ligandbinding domain, a transmembrane domain and a

cytoplasmic serine-threonine kinase domain. This gene may be associated with susceptibility to preeclampsia, a pregnancy-related disease which can result in maternal and fetal morbidity and mortality. Alternative splicing results in multiple

> Email: info@dimabio.com Website: www.dimabio.com

transcript variants of this gene. [provided by RefSeq, Jun 2013]

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)






Figure 1. Human ACVR2A Protein, hFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

