

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	ADA2C
Synonyms	ADRA2L2, ADRA2RL2, ADRARL2, ALPHA2CAR
Description	Human ADA2C-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P18825
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome, GPCRDB Class A Rhodopsin-like,Monoamine GPCRs,
Protein Pathways	
Molecular Weight	The human full length ADA2C-Strep protein has a MW of 49.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Alpha-2-adrenergic receptors are members of the G protein-coupled receptor superfamily. They include 3 highly homologous subtypes: alpha2A, alpha2B, and alpha2C. These receptors have a critical role in regulating neurotransmitter release from sympathetic nerves and from adrenergic neurons in the central nervous system. The mouse studies revealed that both the alpha2A and alpha2C subtypes were required for normal presynaptic control of transmitter release from sympathetic nerves in the heart and from central noradrenergic neurons. The alpha2A subtype inhibited transmitter release at high stimulation frequencies, whereas the alpha2C subtype modulated neurotransmission at lower levels of nerve activity. This gene encodes the alpha2C subtype, which contains no introns in either its coding or untranslated sequences. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

