

## **PRODUCT INFORMATION**

**ANPEP Target** 

**Synonyms** APN;CD13;GP150;LAP1;P150;PEPN

Recombinant human ANPEP protein with C-Description

terminal 6×His tag

**Delivery** In Stock **Uniprot ID** P15144 **Expression Host** HFK293 C-6×His Tag Tag

**Molecular Weight** 

Storage & Shipping

Background

Molecular ANPEP(Lys69-Lys967) 6×His tag Characterization

The protein has a predicted molecular mass of

103.6 kDa after removal of the signal peptide. The apparent molecular mass of ANPEP-His is approximately 130-250 kDa due to glycosylation. The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in other plasma membranes. In the small intestine aminopeptidase N plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Its function in proximal tubular epithelial cells and other cell types is less clear. The large extracellular carboxyterminal domain contains a pentapeptide consénsus sequence characteristic of members of the zinc-binding metalloproteinase superfamily. Sequence comparisons with known enzymes of this class showed that CD13 and aminopeptidase N are

identical. The latter enzyme was thought to be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the CNS. This membrane-bound zinc metalloprotease is known to serve as a receptor for the HCoV-229E alphacoronavirus as well as other

non-human coronaviruses. This gene has also been shown to promote angiogenesis, tumor growth, and metastasis and defects in this gene are associated with various types of leukemia and lymphoma. [provided by RefSeq, Apr 2020]

Email: info@dimabio.com

Website: www.dimabio.com

**Usage** Research use only Unconjugated Conjugate

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Human ANPEP Protein, His Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

