

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	AQP2
Synonyms	AQP-CD; NDI2; WCH-CD
Description	Human AQP2 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P41181
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length AQP2 protein has a MW of 28.2 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Formulation & Reconstitution	A water channel protein located in the kidney collecting tubule. It belongs to the MIP/aquaporin family, some members of which are clustered together on chromosome 12q13. Mutations in this gene have been linked to autosomal dominant and recessive forms of nephrogenic diabetes insipidus.
Storage&Shipping	
Background	
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate AQP2-Nanodisc
0.2 μ g Human AQP2-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag Tag AQP2-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with AQP2-nanodisc is 4.177ng/ml.




Figure 2. Human AQP2-Nanodisc, Flag Tag on SDS-PAGE

