

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	AQP7
Synonyms	AQP7L; AQPap; GLYCQTL
Description	Human AQP7-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	O14520
Expression Host	HEK293
Protein Families	Druggable Genome, Transmembrane
Protein Pathways	PPAR signaling pathway
Molecular Weight	The human full length AQP7-Strep protein has a MW of 37.2 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Formulation & Reconstitution	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage&Shipping	A member of the aquaporin family of water-selective membrane channels. The encoded protein localizes to the plasma membrane and allows movement of water, glycerol and urea across cell membranes. This gene is highly expressed in the adipose tissue where the encoded protein facilitates efflux of glycerol. In the proximal straight tubules of kidney, the encoded protein is localized to the apical membrane and prevents excretion of glycerol into urine. The encoded protein is present in spermatids, as well as in the testicular and epididymal spermatozoa suggesting an important role in late spermatogenesis. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms.
Background	Research use only
Usage	Unconjugated



ELISA assay to evaluate AQP7-Strep-Nanodisc
0.2 μ g Human AQP7-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag AQP7-Strep-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with AQP7-Strep-nanodisc is 5.153ng/ml.




Figure 2. Human AQP7-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

