

## PRODUCT INFORMATION

<b>Target</b>	B7-H7
<b>Synonyms</b>	B7y; B7H7; B7-H5; HHLA2
<b>Description</b>	Recombinant human B7-H7(132-234) Protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9UM44
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc tag
<b>Molecular Characterization</b>	B7-H7(Asn132-Met234) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 37.9 kDa after removal of the signal peptide. The apparent molecular mass of B7-H7(132-234)-hFc is approximately 35-55 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a protein ligand found on the surface of monocytes. The encoded protein is thought to regulate cell-mediated immunity by binding to a receptor on T lymphocytes and inhibiting the proliferation of these cells. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human B7-H7(132-234) Protein, hFc Tag on SDS-PAGE under reducing condition.

