

PRODUCT INFORMATION

Target	BCL2L1
Synonyms	Bcl-X;BCL-XL/S;BCL2L;BCLX;PPP1R52
Description	Recombinant human BCL2L1 protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q07817
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	BCL2L1(Met1-Arg212) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 49.9 kDa after removal of the signal peptide. The apparent molecular mass of BCL2L1-hFc is approximately 55-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an apoptotic activator. [provided by RefSeq, Dec 2015]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human BCL2L1 Protein, hFc Tag on SDS-PAGE under reducing condition.

