

PRODUCT INFORMATION

| | |
|---|--|
| Target | BTK |
| Synonyms | AGMX1;AT;ATK;BPK;IGHD3;IMD1;PSCTK1;XLA |
| Description | Recombinant Human BTK Protein with C-terminal 3×Flag tag |
| Delivery | In Stock |
| Uniprot ID | Q06187 |
| Expression Host | HEK293 |
| Tag | C-3×Flag Tag |
| Molecular Characterization | BTK(Met1-Ser659) 3×Flag tag |
| Molecular Weight | The protein has a predicted molecular mass of 79.2 kDa after removal of the signal peptide. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2013] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human BTK Protein, Flag Tag on SDS-PAGE under reducing condition.

