

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	C5AR1
Synonyms	C5A; C5AR; C5R1; CD88
Description	Human C5AR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P21730
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Complement and coagulation cascades, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length C5AR1 protein has a MW of 39.3 kDa
Background	Receptor for the chemotactic and inflammatory peptide anaphylatoxin C5a. The ligand interacts with at least two sites on the receptor: a high-affinity site on the extracellular N-terminus, and a second site in the transmembrane region which activates downstream signaling events. Receptor activation stimulates chemotaxis, granule enzyme release, intracellular calcium release and superoxide anion production
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C(Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate C5AR1-Nanodisc
0.2 μ g Human C5AR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag C5AR1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with C5AR1-Nanodisc is 4.088ng/ml.



Figure2. Human C5AR1-Nanodisc, Flag Tag on SDS-PAGE

