

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	CAC1C
Synonyms	CACH2, CACN2, CACNL1A1, CCHL1A1, CaV1.2, LQT8, TS, TS. LQT8
Description	Human CAC1C-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q13936
Expression Host	HEK293
Protein Families	Ion Channels: Calcium
Protein Pathways	N/A
Molecular Weight	The human full length CAC1C-Strep protein has a MW of 249 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes an alpha-1 subunit of a voltage-dependent calcium channel. Calcium channels mediate the influx of calcium ions into the cell upon membrane polarization. The alpha-1 subunit consists of 24 transmembrane segments and forms the pore through which ions pass into the cell. The calcium channel consists of a complex of alpha-1, alpha-2/delta, beta, and gamma subunits in a 1:1:1:1 ratio. There are multiple isoforms of each of these proteins, either encoded by different genes or the result of alternative splicing of transcripts. The protein encoded by this gene binds to and is inhibited by dihydropyridine. Alternative splicing results in many transcript variants encoding different proteins. Some of the predicted proteins may not produce functional ion channel subunits. [provided by RefSeq, Oct 2012]
Usage	Research use only
Conjugate	Unconjugated

